

Combining Cytotoxic and Targeted Therapies for Lung Cancer

Matjaz Zwitter, MD, PhD

In this issue, Heist et al.¹ present experience from a phase I to II trial of a combination of topotecan and targeted drug AT-101 for patients with small cell lung cancer in progression after first-line chemotherapy. Although the trial did not include a control group and the study population was relatively small, it nevertheless seems that this combination of a targeted drug and chemotherapy is not superior to chemotherapy alone.

A similar conclusion may be derived from five other studies on advanced non-small cell lung cancer with a total of more than 5000 patients. Two trials (INTACT 1 and INTACT 2) tested chemotherapy alone or in combination with gefitinib^{2,3} and the other two (TRIBUTE and TALENT) tested chemotherapy with or without erlotinib.^{4,5} In all four trials, adding a tyrosine kinase inhibitor to chemotherapy did not prolong the time to progression or overall survival. The fifth study is the recently published FLEX trial in which chemotherapy was given with or without cetuximab.⁶ Addition of cetuximab led to moderate yet statistically significant prolongation of overall survival. However, there was an important difference in the duration of the treatment between the two arms: patients in the chemotherapy-only arm stopped all anticancer treatment after a maximum of 4 months, whereas the cetuximab arm continued with the targeted drug until progression. A close look at the data reveals overlapping survival curves during the first 6 months and their separation only after 8 months of treatment. Therefore, it remains uncertain whether the combination is truly superior to chemotherapy alone. It may well be that the advantage of the cetuximab arm in the FLEX trial was due to the benefit of prolongation of the primary treatment. In other words, it is not the superiority of the combination but longer duration of primary treatment that led to better survival—a phenomenon seen also in positive trials of maintenance or early second-line treatment with gemcitabine, pemetrexed, or erlotinib.^{7–9}

After negative experience from these six trials, should we continue with efforts to combine chemotherapy and targeted drugs or should we abandon any further research in this direction? To approach this question, we have to see what all six trials have in common and try to answer why they failed. All six trials are similar in one parameter: they applied chemotherapy and targeted drugs simultaneously. Schedules that apply chemotherapy and targeted drugs simultaneously ignore one crucial aspect: potential antagonism between the two classes of drugs. Sensitivity to targeted drugs may well push tumor cells to the dormant phases of the mitotic cycle and render them resistant to classic cytotoxic agents. If this is true, then negative results from simultaneous use of cytotoxic drugs and targeted agents do not come as a surprise.

Time separation between chemotherapy and targeted drugs is an approach that avoids the aforementioned antagonism between the two classes of drugs. An additional advantage of such an intermittent schedule is prevention of tumor repopulation that occurs during the gap between individual applications of chemotherapy.¹⁰ In vitro research has clearly demonstrated that schedules with different timing of chemotherapy and of targeted drugs lead to their synergistic or antagonistic antitumor activity.^{11–13} Preliminary promising experience from clinical trials conducted in United States,¹⁴ Hong Kong,¹⁵ and

Institute of Oncology, Ljubljana, Slovenia.

Disclosure: The author declares no conflict of interest.

Address for correspondence: Matjaz Zwitter, MD, PhD, Institute of Oncology, Zaloska 2, 1000 Ljubljana, Slovenia. E-mail: mzwitter@onko-i.si

Copyright © 2010 by the International Association for the Study of Lung Cancer

ISSN: 1556-0864/10/0510-1498

Slovenia (M. Zwitter, unpublished data) supports the concept of intermittent schedules. Deeper understanding of general tumor biology, the characteristics of each individual tumor, and optimal timing of our interventions should lead to fewer disappointments in future clinical research.

REFERENCES

1. Heist R, Fain J, Chinnasami B, et al. Phase I/II study of AT-101 with topotecan in relapsed and refractory small cell lung cancer (SCLC). *J Thorac Oncol* 2010;5:1637–1643.
2. Giaccone G, Herbst RS, Manegold C, et al. Gefitinib in combination with gemcitabine and cisplatin in advanced non-small-cell lung cancer: a phase III trial—INTACT 1. *J Clin Oncol* 2004;22:777–784.
3. Herbst RS, Giaccone G, Schiller JH, et al. Gefitinib in combination with paclitaxel and carboplatin in advanced non-small-cell lung cancer: a phase III trial—INTACT 2. *J Clin Oncol* 2004;22:785–794.
4. Herbst RS, Prager D, Herman R, et al. TRIBUTE: a phase III trial of erlotinib hydrochloride (OSI-774) combined with carboplatin and paclitaxel chemotherapy in advanced non-small-cell lung cancer. *J Clin Oncol* 2005;23:5892–5899.
5. Gatzemeier U, Pluzanska A, Szczesna A, et al. Phase III study of erlotinib in combination with cisplatin and gemcitabine in advanced non-small-cell lung cancer: the Tarceva Lung Cancer Investigation Trial. *J Clin Oncol* 2007;25:1545–1552.
6. Pirker R, Pereira JR, Szczesna A, et al. Cetuximab plus chemotherapy in patients with advanced non-small-cell lung cancer (FLEX): an open-label randomised phase III trial. *Lancet* 2009;373:1525–1531.
7. Brodowicz T, Krzakowski M, Zwitter M, et al. Cisplatin and gemcitabine first-line chemotherapy followed by maintenance gemcitabine or best supportive care in advanced non-small cell lung cancer: a phase III trial. *Lung Cancer* 2006;52:155–163.
8. Ciuleanu T, Brodowicz T, Zielinski C, et al. Maintenance pemetrexed plus best supportive care versus placebo plus best supportive care for non-small-cell lung cancer: a randomised, double-blind, phase 3 study. *Lancet* 2009;374:1432–1440.
9. Mok TS, Ramalingam SS. Maintenance therapy in nonsmall-cell lung cancer: a new treatment paradigm. *Cancer* 2009;115:5143–5154.
10. Davis AJ, Tannock JF. Repopulation of tumour cells between cycles of chemotherapy: a neglected factor. *Lancet Oncol* 2000;1:86–93.
11. Mahaffey CM, Davies AM, Lara PN Jr, et al. Schedule-dependent apoptosis in K-ras mutant non-small-cell lung cancer cell lines treated with docetaxel and erlotinib: rationale for pharmacodynamic separation. *Clin Lung Cancer* 2007;8:548–553.
12. Li T, Ling YH, Goldman ID, et al. Schedule-dependent cytotoxic synergism of pemetrexed and erlotinib in human non-small cell lung cancer cells. *Clin Cancer Res* 2007;13:3413–3422.
13. Solit DB, She Y, Lobo J, et al. Pulsatile administration of the epidermal growth factor receptor inhibitor gefitinib is significantly more effective than continuous dosing for sensitizing tumors to paclitaxel. *Clin Cancer Res* 2005;11:1983–1989.
14. Davies AM, Ho C, Beckett L, et al. Intermittent erlotinib in combination with pemetrexed: phase I schedules designed to achieve pharmacodynamic separation. *J Thorac Oncol* 2009;4:862–868.
15. Mok TS, Wu YL, Yu CJ, et al. Randomized, placebo-controlled, phase II study of sequential erlotinib and chemotherapy as first-line treatment for advanced non-small-cell lung cancer. *J Clin Oncol* 2009;27:5080–5087.